

National Library
of Medicine

PubMed

PubMed	Nucleotide	Protein	Genome	Structure	PopSet
Search PubMed	for				Go Clear
Limits		Preview/Index		History	Clipboard

Entrez PubMed

☐ 1: *Adv Exp Med Biol* 1992;321:1-5

Related Articles, Books

PubMed Services

Pancreatic islet cell regeneration and growth. Introduction.**Vinik AI**

Eastern Virginia Medical School, Diabetes Institutes, Norfolk 23510.

Publication Types:

- Review
- Review, tutorial

Related Resources

PMID: 1449076, UI: 93080098

Display	Abstract	<input checked="" type="checkbox"/>	Save	Text	Order	Add to Clipboard
---------	----------	-------------------------------------	------	------	-------	------------------

Write to the Help DeskNCBI | NLM | NIHDepartment of Health & Human ServicesFreedom of Information Act | Disclaimer

INTRODUCTION

Aaron I. Vinik, M.D., Ph.D.¹

¹Eastern Virginia Medical School
The Diabetes Institutes
Norfolk, Virginia 23510

This symposium, held in June 1991, was a gathering of international scientists to exchange their views on current concepts of cell growth and differentiation. Each scientist was asked to present a topic of their research related to cell growth and regeneration and to participate in a round table conference elaborating on current knowledge and sharing their experiences. By furthering this promising area of endeavor, a means of understanding ontogeny of cell development and of providing insights into tumor biology would prevail. Of prime importance, was the anticipation that new information from a better understanding of the normal evolution of the pancreatic islet would generate alternative approaches to curing diabetes. This forward serves as a short introduction to the concept of pancreatic islet regeneration and the models currently in use to study the process.

DEVELOPMENTAL ORIGIN OF ISLETS DURING EMBRYOGENESIS

The developing pancreas appears as a protrusion from the dorsal surface of the embryonic gut.¹ The different islet cell types appear sequentially during development *in vivo*. It therefore seems reasonable to propose that coordinated growth is dependent upon specificity of growth factors.

Islet cells *in vivo* also express several neuroectodermal antigens, for example, PGP 9.5,² neurone-specific enolase (NSE),³ synaptophysin,⁴ A2B5,⁵ phenylethanolamine N-methyltransferase (PNMT) and aromatic amino acid decarboxylase (AADC).⁶ The endocrine cells of the GEP axis are capable of amine precursor uptake and decarboxylation and have, therefore, been given the acronym APUD.⁷ The morphologic similarity of APUD cells suggested a common embryologic origin, which was believed to be the neural crest but later revised to include the neuroectoderm, or in the case of some of the endocrine cells, from the dorsal placoderm.

Studies by Ledouarin,⁸ Pictet,⁹ Andrew,¹⁰ and their coworkers have cast doubt on this hypothesis, and most workers agree that these cells should be classified according to their secretory products, i.e., gastrin, somatostatin, glucagon, PP, etc. However, it is now thought that β -cells do not have APUD characteristics and are likely to be derived from gut, but express neuronal antigens such as the catecholamine biosynthetic enzyme tyrosine hydroxylase (TH).¹¹ The generally held belief that the neuronal characteristics of these cells indicated an ectodermal origin during mammalian embryogenesis has largely been dispelled.

During development *in vivo*, the phenotype of the mature islet cells appear sequentially. β -cells arise from progenitor cells localized in the pancreatic duct and these precursors transiently express TH while migrating away from the duct to populate a new islet. This suggests that the pancreatic duct is a source of endocrine stem cells throughout embryogenesis without the need to postulate a neuroendocrine origin. This notion is supported by the finding that the embryonic pancreatic duct *in vitro* is able to regenerate a new pancreas containing exocrine and endocrine cells expressing only peptides (mature cells), and cells containing both TH and a hormone (immature cells).^{12,13}

Teitelman has shown that pancreatic cells of endocrine origin can indeed express several neuronal antigens in addition to the peptide hormones.¹¹ She further showed that in the mouse embryo a primitive undifferentiated cell(s) led to sequential appearance of at least 4 different cell types containing either a hormone (e.g., glucagon), a catecholamine enzyme (tyrosine hydroxylase, TH) or combinations of these.⁶ Under appropriate conditions these cells can be shown to differentiate into either neurites or adult endocrine cells. During regeneration, expression of neural antigens by developing cells was found to constitute an early phase to be replaced by the adult hormone secreting counterpart. Rosenberg and Vinik¹⁴ have utilized a model for nesidioblastosis and shown that pancreatic ductal cells are capable of differentiating upon stimulation into adult endocrine cells capable of secreting insulin in a fully regulated manner.

ISLET CELL GROWTH AND DIFFERENTIATION

Factors which control the growth and functional maturation and differentiation of the human endocrine pancreas and gut during the fetal and post-natal periods are incompletely understood. The role of the fetal mesenchyme in epithelial cell development and differentiation appears to be important.¹⁵⁻¹⁷ Possible mechanisms of action include: (i) secretion of an inducing or transforming hormonal growth factor, (ii) information exchange through cell-to-cell contact via paracrine and juxtacrine actions of locally elaborated growth factors, and (iii) production of an extracellular matrix rich in growth promoting factors. The soluble peptide growth factors are trophic substances that regulate both cell proliferation and differentiation and may be linked to islet growth.

One family of growth factors that may be implicated in islet growth are the somatomedins and their binding proteins. Insulin-like growth factors (IGFs) are important mediators of fetal and postnatal growth. Whereas these growth factors circulate (attached to binding proteins) they also act locally. Fetal rat islets release both IGF-I and IGF-II *in vitro* which may contribute to growth hormone-induced DNA synthesis.¹⁸⁻²¹ It is apparent that the role of IGF-I, and the binding proteins, especially in the adult pancreas is far from clear. In this symposium Drs. LeRoith and Lauterio discuss aspects of IGF physiology, and LeRoith focuses upon differences in the role of the IGF's in regeneration of adult and fetal tissues. Dr. Hill emphasizes the role of the IGF's and their binding proteins in islet regeneration in the fetal pancreas at a time when the pancreas is susceptible to the influence of this particular group of growth factors. Dr. Bonner Weir also points out that in their model of islet regeneration after 90% pancreatectomy there is enhanced IGF gene expression in the ductules and certain connective tissue cells in contrast with the normal expression in capillary endothelial cells suggesting that IGFs may participate in the regeneration process after pancreatectomy in the rat.²² Dr. Nielsen, however, contests the suggestion that the IGF's are important as pancreatic trophic factors and based upon his observations growth hormone itself may be pertinent, at least in pregnancy, a state in which islet hypertrophy is found.

Several important glycoprotein components of extracellular matrix - the "integrins" - have also been recognized as playing a role in cell growth and differentiation. For example, fibronectin, laminin and tenascin.^{23,24} Dr. Le Beau elaborates upon the role the integrins may play in cell regeneration. The role of these factors in the maintenance and replacement of a functional islet cell mass in the adult pancreas remains to be determined.

ISLET CELL PROLIFERATION IN THE POST-NATAL PERIOD

Several models designed to induce exocrine and endocrine pancreatic regeneration have been developed. In the model developed by Bonner-Weir and colleagues,^{22,25} there is regeneration of both exocrine and endocrine tissue following a 90% pancreatectomy in which the increase in β -cell mass occurs as a result of the replication of existing β -cells and not necessarily because of a process of new islet formation.²² This group have reported increased IGF-I mRNA production by capillary endothelial cells and proliferating ductules which may contribute to both endocrine and exocrine pancreas regeneration, but the precise

role of IGFs needs to be has not been excluded.²²

Terazono et al developed a model in which the pancreas was treated with either methylauriothioglucose resulting in a substance termed *reg* gene.²⁸ Human *reg* mRNA levels in gastric mucosa expressed ectopically in transformed, proliferative *reg* is expressed in acinar growth factor. Dr. O. regeneration, elaborates this growth factor in islet and Rafacloff.

Expression of an islet tissue.³⁰ The significance of regeneration studied by over a period of weeks to

A second version of colleagues,³¹ deals with Hospital (NEDH) rats to demonstrate that tumor with a profound reduction that results in rapid β -mRNA levels. Whether high levels of *reg* mRNA ongoing at a low, constant this gene product.

Another model has been observed in transgenic mice. These mice, in which duct cell proliferation along the apical region indicating that diabetes destructive process. To induce regeneration may

In 1982, we developed 8 weeks of age are fertile of cell proliferation the hamster pancreatic duct was observed. The proliferation and differentiation processes were shown to an extract prepared from other hamsters, but this pancreas was administered developments in the use called flitropin.

The presence of the has been hypothesized but reviews the current knowledge carried out to further determine the release of a variety and differentiation of with factors important in possible mechanism of also discussed.

in can indeed express several
ie further showed that in the
ntial appearance of at least 4
n), a catecholamine enzyme
appropriate conditions these
ult endocrine cells. During
ls was found to constitute an
counterpart. Rosenberg and
rat pancreatic ductal cells are
ne cells capable of secreting

ion and differentiation of the
atal periods are incompletely
elial cell development and
nisms of action include: (i)
or, (ii) information exchange
; of locally elaborated growth
rowth promoting factors. The
ate both cell proliferation and

ed in islet growth are the
factors (IGFs) are important
factors circulate (attached to
oth IGF-I and IGF-II *in vitro*
esis.¹⁸⁻²¹ It is apparent that
ult pancreas is far from clear.
acts of IGF physiology, and
egeneration of adult and fetal
ir binding proteins in islet
is susceptible to the influence
also points out that in their
hanced IGF gene expression
with the normal expression in
te in the regeneration process
nests the suggestion that the
pon his observations growth
in which islet hypertrophy is

ular matrix - the "integrins"-
differentiation. For example,
es upon the role the integrins
maintenance and replacement
e determined.

PERIOD

e pancreatic regeneration have
and colleagues,^{22,25} there is
ng a 90% pancreatectomy in
lication of existing β -cells and
²² This group have reported
ells and proliferating ductules
s regeneration, but the precise

role of IGFs needs to be elucidated and the possibility that other growth factors participate
has not been excluded.²²

Terazono et al developed a model in which 90% pancreatectomized rats or mice are
treated with either nicotinamide, (a poly ADP-ribose synthetase inhibitor), or
aurothioglucose resulting in exocrine and endocrine cell regeneration, and the appearance of
a substance termed *reg* protein.^{26,27} The gene encoding this protein has been termed the *reg*
gene.²⁸ Human *reg* mRNA has been detected predominantly in the pancreas, and at lower
levels in gastric mucosa and in the kidney.²⁹ *Reg* gene protein has also been found to be
expressed ectopically in colon and rectal tumors,²⁹ linking enhanced *reg* expression to the
transformed, proliferative state, at least for some cell types. Current evidence suggests that
reg is expressed in acinar tissue and not regenerating islets. It may therefore be a paracrine
growth factor. Dr. Okamoto, who pioneered the work on *Reg* gene and pancreatic
regeneration, elaborates here upon his new findings. The controversy over the relevance of
this growth factor in islet regeneration in other models is further discussed by Drs. Newgard
and Rafaeloff.

Expression of an homologous gene, termed *rig*, has been identified in insulinoma
tissue.³⁰ The significance of these genes remains to be determined. In both models of
regeneration studied by Okamoto and colleagues,^{26,27,29} islet size increased above normal
over a period of weeks to months.

A second version of the 90% pancreatectomized rat model studied by Newgard and
colleagues,³¹ deals with *reg* expression in insulinoma-bearing New England Deaconess
Hospital (NEDH) rats relative to normal controls and following tumor resection. They
demonstrate that tumor implantation causes a sharp reduction in *reg* expression associated
with a profound reduction in non-tumor islet size and that removal of the tumor, a maneuver
that results in rapid β -cell proliferation, results in a large but transient induction in *reg*
mRNA levels. Whether *reg* protein is β -cytotropic is still an open question. The fact that
high levels of *reg* mRNA are present in normal animals, in which β -cell replication is
ongoing at a low, constitutive rate, are seemingly at odds with a growth-promoting role for
this gene product.

Another model has been suggested by Dr. Sarvetnick³² in which a regenerative process
is observed in transgenic mice expressing Interferon-gamma in their pancreatic β -cells.
These mice, in which diabetes ensues following immunodestruction of their β -cells, show
duct cell proliferation and the appearance of more primitive neuroendocrine progenitor cells
along the apical regions of the ducts. Dr. Sarvetnick discusses her most recent findings
indicating that diabetes may not develop provided that the regenerative process outstrips the
destructive process. This, we believe, is an important principle whereby an approach to
induce regeneration may not be unreasonable in our quest for a cure for diabetes.

In 1982, we developed a unique model for islet regeneration in hamsters.^{33,34} Hamsters
8 weeks of age are fertile, considered to be adult animals and respond better to the induction
of cell proliferation than do older animals.^{33,34} By producing partial obstruction of the
hamster pancreatic duct by cellophane wrapping, new islet formation from ductal elements
was observed. The mechanism by which partial obstruction in our model induces cell
proliferation and differentiation is unknown. Using a parabiotic experimental design, these
processes were shown to be mediated by paracrine and/or autocrine mechanisms.³⁰ Indeed,
an extract prepared from a wrapped pancreas exhibited trophic activity when injected into
other hamsters, but this was not observed when an extract prepared from a non-wrapped
pancreas was administered. Drs. Rosenberg and Vinik review the data and the more recent
developments in the use of a cytosol extract, containing the growth factor which we have
called Ilotropin.

The presence of this specific growth factor, Ilotropin, in the β -cell cytosol extract has
been hypothesized but the identity of the peptide had not been established. Dr. Pittenger
reviews the current knowledge of the nature of the growth factor and the studies he has
carried out to further define its characteristics. However, cellophane wrapping may initiate
the release of a variety of growth factors that contribute to the coordinated, timely growth
and differentiation of ductal cells. These possibilities and other factors that may be shared
with factors important in the nervous system are highlighted by members of the faculty. The
possible mechanism of action of the various factors as well as therapeutic applications are
also discussed.

GROWTH FACTOR(S) AND NEOPLASIA

A great deal of interest is now being focused upon the factors responsible for initiation of growth, increase in cell number and size, differentiation into adult endocrine cells, growth cessation and cell maintenance.^{36,37} The coincidental findings that the multiple endocrine neoplasia, type 1 syndrome (MEN-1) (combined occurrence of tumors of the pituitary, pancreas and parathyroid glands) is associated with the loss of alleles on chromosome 11,^{38,39} the same chromosome on which the insulin gene has been located;⁴⁰ the finding of parathyroid mitogenic activity in the plasma of patients with MEN-1;^{36,37} and evidence that patients with MEN-1 might also secrete mitogenic factors for pancreatic islet-cells into plasma,⁴¹ suggests a role for genetically determined circulating growth factors in the 'growth initiation' of these tumors.

It is apparent, therefore, that the conference was timely and that the information pertinent to our appreciation of cell growth and differentiation is relevant to a possible cure for diabetes as well as a clearer understanding of factors that may be involved in pancreatic tumor formation. Conceptually, these conditions represent the two ends of a spectrum - diabetes with endocrine cell failure and endocrine neoplasia with unbridled cell growth.

REFERENCES

1. R.L. Pictet and J.W. Rutter, Development of the embryonic endocrine pancreas, in: "Handbook of Physiology," D.F. Steiner and M. Frenkel, eds., Washington D.C., American Physiological Society (1972).
2. K.J. Thompson, J.F. Doran, P. Jackson, A.P. Dhillon, and J. Rode, PGP 9.5: a new marker for vertebrate neurons and endocrine cells, *Brain Res.* 278:224-28 (1983).
3. Polak, et al. Neuron-specific enolase, a marker for neuro-endocrine cells, in: *Evolution and Tumor Pathology of the Neuro-endocrine System*. Elsevier, New York (1984).
4. B. Weidenmann, W.W. Franke, C. Kuhn, R. Moll, and V.E. Gould, Synaptophysin: a marker protein for neuro-endocrine cells and neoplasms, *Proc Natl Acad Sci USA.* 83:3500-04 (1986).
5. G.S. Eisenbarth, K. Shimizu, M.A. Bowring, and S. Wells, Expression of receptors for fetanin toxin and monoclonal antibody A2B2 by pancreatic islet cells, *Proc Natl Acad Sci USA.* 79:5066-70 (1982).
6. G. Teitelman, T.H. Joh, and D.J. Reis, Transformation of catecholaminergic precursor into glucagon (A) cells in mouse embryonic pancreas, *Proc Natl Acad Sci USA.* 78:5225-29 (1991).
7. A.G.E. Pearse, Common cytochemical and ultrastructural characteristics of cells producing polypeptide hormones (the APUD series) and their relevance to thyroid and ultimobronchial C cells and calcitonin, *Proc R Soc Lond (Biol).* 170:71 (1968).
8. N.M. LeDourian and M.A. Teillet, The migration of neural crest cells to the wall of the digestive tract in avian embryo, *J Embryol Exp Morphol.* 30:31 (1973).
9. R.L. Pictet, L.B. Rall, P. Phelps, and W.J. Rutter, The neural crest and the origin of the insulin-producing and other gastrointestinal hormone-producing cells, *Science.* 191:191 (1967).
10. A. Andrew, An experimental investigation into the possible neural crest origin of pancreatic APUD (islet) cells, *J Embryol Exp Morphol.* 35:577 (1976).
11. G. Teitelman and J.K. Lee, Cell lineage analysis of pancreatic islet cell development: Glucagon and insulin cells arise from catecholaminergic precursor present in the pancreatic duct, *Devel Biol.* 121:454-56 (1987).
12. G. Teitelman, J. Lee, and D.J. Reis, Differentiation of prospective mouse pancreatic islet cells during development *in vitro* and during regeneration, *Dev Biol.* 120:425-33 (1987).
13. R.W. Dudek and I.E. Lawrence, Morphologic evidence of interaction between adult ductal epithelium of pancreas and fetal foregut mesenchyme, *Diabetes.* 37:891-900 (1988).
14. L. Rosenberg, W.P. Duguid, and A.I. Vinik, Cell proliferation in the pancreas of the Syrian golden hamster, *Dig Dis Sci.* 32:1185 (1987).
15. R.L. Pictet, L. Rall, M. de Gasparo, and W.J. Rutter, Regulation of differentiation of endocrine cells during pancreatic development *in-vitro*, in: "Early diabetes in early life," R.A. Camerini-Davalos and H.S. Cole, eds., Academic Press, New York (1975).
16. B.S. Spooner, H.I. Cohen, and J. Faubion, Development of the embryonic mammalian pancreas: The relationship between morphogenesis and cytodifferentiation, *Dev Biol.* 61:119 (1977).
17. R. Montesano, P. Mouron, M. Amherdt, and L. Orci, Collagen matrix promotes reorganization of pancreatic endocrine cell monolayers into islet-like organoids, *J Cell Biol.* 97:935 (1983).
18. I. Sweeney and D.J. Hill, Growth hormone regulation of DNA replication, but not insulin production, is partly mediated by somatomedin C/insulin-like growth factor-I in isolated pancreatic islets from adult rats, *Diabetologia.* 32:191-97 (1989).
19. D.J. Hill, A. Frazer, I. Swe, P.K. Wirtham, and R.D.G. Milner, Somatostatin-C in human fetal pancreas, *Diabetes.* 36:465 (1987).
20. A. Rabinovitch, C. Quigley, et al., Insulin-like growth factor activity (an insulin-like monolayer cultures, *J. Clin. Endocrinol.* 61:1185 (1985).
21. J.M. Bryson, B.E. Tuch, et al., Insulin-like growth factor activity in the pancreas in culture, *J. Clin. Endocrinol.* 61:1185 (1985).
22. J.S. Brockenbrough, G.C. Brockenbrough, et al., Insulin-like growth factor activity in the pancreas after 90% pancreatectomy, *J. Clin. Endocrinol.* 61:1185 (1985).
23. M.J. Politis, Exogenous insulin-like growth factor activity in the pancreas after reconstruction, *Reconstr Surg.* 83:22 (1985).
24. R. Chiquet-Ehrismann, B. Chiquet, et al., Insulin-like growth factor protein involved in tissue repair, *J. Clin. Endocrinol.* 61:1185 (1985).
25. F. Smith, K. Rosen, L. Vitiello, et al., Insulin-like growth factor activity is localized to capillaries in the pancreas, *J. Clin. Endocrinol.* 61:1185 (1985).
26. K. Terazono, H. Yamamoto, et al., Insulin-like growth factor gene activated in regenerating pancreas, *J. Clin. Endocrinol.* 61:1185 (1985).
27. K. Terazono, Y. Uchiyama, et al., Expression of reg protein in regenerating pancreatic secretory granules, *J. Clin. Endocrinol.* 61:1185 (1985).
28. T. Watanabe, H. Yonekura, et al., Expression of human reg gene in the pancreas, *J. Clin. Endocrinol.* 61:1185 (1985).
29. C. Miyauchi, et al., Expression of reg protein in the pancreas, *J. Clin. Endocrinol.* 61:1185 (1985).
30. S. Takasawa, K. Yamamoto, et al., Insulin-like growth factor gene activated in regenerating pancreas, *J. Clin. Endocrinol.* 61:1185 (1985).
31. L. Chen, M. Appel, T. Alizadeh, et al., Regulating Islet Regeneration, *J. Clin. Endocrinol.* 61:1185 (1985).
32. N. Sarvetnick, Islet cell death, *J. Clin. Endocrinol.* 61:1185 (1985).
33. L. Rosenberg, W.P. Duguid, et al., Diabetes in the Syrian hamster, *J. Clin. Endocrinol.* 61:1185 (1985).
34. L. Rosenberg and A.I. Vinik, Paracrine and/or autocrine regulation of islet cell growth, *J. Clin. Endocrinol.* 61:1185 (1985).
35. L. Rosenberg and A.I. Vinik, Islet cell growth from the "wrapped" islet, *J. Clin. Endocrinol.* 61:1185 (1985).
36. M.L. Brandt, G.D. Aurbach, et al., Familial multiple endocrine neoplasia, *J. Clin. Endocrinol.* 61:1185 (1985).
37. S.J. Marx, K. Sakaguchi, et al., Insulin-like growth factor in plasma from multiple endocrine neoplasia, *J. Clin. Endocrinol.* 61:1185 (1985).
38. C. Larsson, B. Skogseid, et al., Insulin-like growth factor I gene maps to chromosome 11, *J. Clin. Endocrinol.* 61:1185 (1985).
39. R.V. Thakker, P. Bouloux, et al., Multiple endocrine neoplasia, *J. Clin. Endocrinol.* 61:1185 (1985).
40. D. Owerbach, G.I. Bell, et al., Insulin-like growth factor I gene maps to chromosome 11, *J. Clin. Endocrinol.* 61:1185 (1985).
41. M.K. McLeod, A.M. Tuttle, et al., Insulin-like growth factor I gene maps to chromosome 11, *J. Clin. Endocrinol.* 61:1185 (1985).

ctors responsible for initiation of adult endocrine cells, growth factors that the multiple endocrine tumors of the pituitary, loss of alleles on chromosome 11 have been located;⁴⁰ the finding of MEN-1;^{36,37} and evidence that for pancreatic islet-cells into growth factors in the 'growth

ous and that the information is relevant to a possible cure may be involved in pancreatic the two ends of a spectrum - with unbridled cell growth.

pancreas, in: "Handbook of American Physiological Society

P 9.5: a new marker for vertebrate

cells, in: Evolution and Tumor (1984).

pancreatic polypeptide: a marker protein for 3:3500-04 (1986).

of receptors for fetanin toxin and J Biol Chem. 257:5066-70 (1982).

precursor into glucagon (A) 5225-29 (1991).

cells of cells producing polypeptide timobronchial C cells and calcitonin,

to the wall of the digestive tract in

the origin of the insulin-producing 191 (1967).

st origin of pancreatic APUD (islet)

development: Glucagon and pancreatic duct, Devel Biol.

use pancreatic islet cells during 33 (1987).

between adult ductal epithelium of 1988).

pancreas of the Syrian golden

differentiation of endocrine cells during "R.A. Camerini-Davalos and H.S.

unic mammalian pancreas: The J Biol. 61:119 (1977).

promotes reorganization of Cell Biol. 97:935 (1983).

on, but not insulin production, is isolated pancreatic islets from adult

ostatin-C in human fetal pancreas,

20. A. Rabinovitch, C. Quigley, T. Russell, Y. Patel, and D.H. Mintz, Insulin and multiplication stimulating activity (an insulin-like growth factor) stimulate islet β -cell replication in neonatal rat pancreatic monolayer cultures, *Diabetes*. 31:160 (1982).
21. J.M. Bryson, B.E. Tuch, and R.C. Baxter, Production of insulin-like growth factor-II by human fetal pancreas in culture, *J Endocrinol*. 121:367-73 (1989).
22. J.S. Brockenbrough, G.C. Weir, and S. Bonner-Weir, Discordance of exocrine and endocrine growth after 90% pancreatectomy in rats, *Diabetes*. 37:232-36 (1988).
23. M.J. Politis, Exogenous laminin induces regenerative changes in traumatized sciatic and optic nerve, *Plas Reconstr Surg*. 83:228-35 (1989).
24. R. Chiquet-Ehrismann, E.J. Mackie, C.A. Pearson, and T. Sakakura, Tenascin: an extracellular matrix protein involved in tissue interactions during fetal development and oncogenesis, *Cell*. 47:131-39 (1986).
25. F. Smith, K. Rosen, L. Villa-Komaroff, et al, Enhanced IGG-I gene expression in regenerating rat pancreas is localized to capillaries and proliferating ductules, *Diabetes*. 39(Suppl 1):66A (1990).
26. K. Terazono, H. Yamamoto, S. Takasawa, K. Shiga, Y. Yonemura, Y. Tochino, and H. Okamoto, A novel gene activated in regenerating islets, *J Biol Chem*. 262:2111 (1988).
27. K. Terazono, Y. Uchiyama, M. Ide, T. Watanabe, H. Yonekura, H. Yamamoto, and H. Okamoto, Expression of *reg* protein in rat regenerating islets and its co-localization with insulin in beta cell secretory granules, *Diabetologia*. 33:1 (1990).
28. T. Watanabe, H. Yonekura, K. Terazono, H. Yamamoto, and H. Okamoto, Complete nucleotide sequence of human *reg* gene and its expression in normal and tumoral tissues, *J Biol Chem*. 265:7432-39 (1990).
29. C. Miyaura, et al, Expression of *reg/PSP*, a pancreatic exocrine gene: Relationship to changes in islet β -cell mass, *Mol Endocrinol*. (1991, in press).
30. S. Takasawa, K. Yamamoto, K. Terazono, and H. Okamoto, Novel gene activated in rat insulinoma, *Diabetes*. 35:1178 (1986).
31. L. Chen, M. Appel, T. Alam, C. Miyaura, A. Sestak, J. O'Neil, R. Unger, and C. Newgard, Factors Regulating Islet Regeneration in the Post-Insulinoma NEDH Rat.
32. N. Sarvetnick, Islet cell destruction and regeneration in IFN- γ transgenic mice, *J Cell Biochem*. CB019:49 (abstract) (1991).
33. L. Rosenberg, W.P. Duguid, R.A. Brown, and A.I. Vinik, Induction of islet cell proliferation will reverse diabetes in the Syrian golden hamster, *Diabetes*. 37:334 (1988).
34. L. Rosenberg and A.I. Vinik, Regulation of pancreatic islet growth and differentiation: Evidence for paracrine and/or autocrine growth factor(s), *Clin Res*. 38:271A (1990).
35. L. Rosenberg and A.I. Vinik, In vitro stimulation of hamster pancreatic duct growth by an extract derived from the "wrapped" pancreas, (submitted for publication).
36. M.L. Brandi, G.D. Aurbach, L.A. Fitzpatrick, et al, Parathyroid mitogenic activity in plasma from patients with familial multiple endocrine neoplasia type 1, *N Engl J Med*. 314:1287-93 (1986).
37. S.J. Marx, K. Sakaguchi, J. Green, G.D. Aurbach, and M.L. Brandi, Mitogenic activity on parathyroid cells in plasma from members of a large kindred with multiple endocrine neoplasia Type 1, *J Clin Endocrinol Metab*. 67:149-53 (1988).
38. C. Larsson, B. Skogseid, K. Oberg, Y. Nakamura, and M. Nordenskjold, Multiple endocrine neoplasia type 1 gene maps to chromosome 11 and is lost in insulinoma, *Nature*. 332:85-87 (1988).
39. R.V. Thakker, P. Bouloux, C. Wooding, K. Chotai, P.M. Broad, N.K. Spurr, G.M. Besser, and J.L. O'Riordan, Association of parathyroid tumors in multiple endocrine neoplasia type 1 with loss of alleles on chromosome 11, *N Engl J Med*. 321:218-24 (1989).
40. D. Owerbach, G.I. Bell, W.J. Rutter, J.A. Brown, and T.B. Shows, The insulin gene is activated on the short arm of chromosome 11 in humans, *Diabetes*. 30:267-70 (1981).
41. M.K. McLeod, A.M. Tuter, N.W. Thompson, and A.I. Vinik, Evidence for a pancreatic islet-cell mitogenic factor in patients with MEN-1, Association for Academic Surgery, Louisville, Kentucky, November 15-18, 1989.